

ENERGY LEVELS AND WAVE FUNCTIONS OF VECTOR BOSONS IN HOMOGENEOUS MAGNETIC FIELD

K.SOGUT, A.HAVARE, I.ACIKGOZ

MERSIN UNIVERSITY, DEPARTMENT OF PHYSICS

(February 1, 2008)

Abstract

We aimed to obtain the energy levels of spin-1 particles moving in a constant magnetic field. The method used here is completely algebraic. In the process to obtain the energy levels the wave function is choosen in terms of Laguerre Polynomials.

I. INTRODUCTION

The problem of finding the states of a particle moving in an external field has been solved for many situations. The relativistic wave equation of spin-1 particles was at first derived by Kemmer in 1939 [1]. This equation is similar to the Dirac equation, but the technique investigated by Kemmer is simpler. Kemmer equation was developed by different matrices.

There are various wave equations to describe the spin-1 particles given in literature [2–5]. The methods used in these equations are quite complicated and give different results when they are compared. The method of Shay and Good is to expand the six-component wave function in terms of complete set of functions. This method leads to an invariant way to take matrix elements, and it allows for arbitrary magnetic dipole moment and electric quadrupole moment of the particle. The eigenvalues of Shay and Good's equation have been obtained by Krase, Lu, Good by using the quite complicated conventional method of solving the differential equation. A general method of obtaining the energy levels of any spin theory

has been proposed by Tsai and Yildiz. They went on to obtain the energy levels of spin-1 theory of Proca and Kemmer and observed that spin-1 theory is consistent only when there is no anomalous -magnetic moment coupling.

The generally accepted method of approach is to define the Lagrange equation of motion and use the differential equation technique to solve for the eigenfunctions and eigenvalues of system. This method becomes complicated when higher spinning particles are handled. In this study we used an algebraic method for obtaining the energy levels of spin-1 particles. This method of approach does not require an explicit solution of the equation. For introducing the method we first used it to obtain the energy levels of spin- $\frac{1}{2}$ particles in section two and by using the eigenfunctions that found for spin- $\frac{1}{2}$ particles moving in terms of Laguerre polynomials in Kemmer equation we obtained the energy levels of spin-1 particles in a homogeneous magnetic field. The vector potential we used is

$$\vec{A} = \frac{1}{2}xB\hat{y} - \frac{1}{2}yB\hat{x}. \quad (1)$$

II. ENERGY LEVELS OF SPIN- $\frac{1}{2}$ PARTICLES

The wave equation is

$$(\gamma^\mu \pi_\mu - m)\Psi(x) = 0 \quad (2)$$

where γ^μ are Dirac matrices and π_μ is electrodynamical four-momentum tensor. The explicit form of this equation is

$$\begin{pmatrix} (E-m) & 0 & -P_z & -\pi_- \\ 0 & (E-m) & -\pi_+ & P_z \\ -P_z & -\pi_- & (E-m) & 0 \\ -\pi_+ & P_z & 0 & (E-m) \end{pmatrix} \begin{pmatrix} \varphi_+ \\ \varphi_- \\ \chi_+ \\ \chi_- \end{pmatrix} = 0 \quad (3)$$

where helicity raising and lowering operators are defined as

$$\pi_+ = \pi_1 + i\pi_2 = e^{i\theta} \left[-i\partial_r + \frac{1}{r}\partial_\theta - \frac{iBre}{2} \right], \quad (4)$$

$$\pi_- = \pi_1 - i\pi_2 = e^{-i\theta} \left[-i\partial_r - \frac{1}{r}\partial_\theta + \frac{iBre}{2} \right]. \quad (5)$$

Before obtaining the energy levels we must note that the effect of helicity raising and lowering operators on Laguerre polynomials.

For $m > 0$

$$\pi_+ L_N^{|m|} F_m = 2i\omega^{\frac{1}{2}} x^{\frac{1}{2}} L_{N-1}^{|m|+1} F_{m+1}, \quad (6)$$

$$\pi_- L_N^{|m|} F_m = -2iN\omega^{\frac{1}{2}} x^{-\frac{1}{2}} L_{N+1}^{|m|-1} F_{m-1}, \quad (7)$$

and for $m < 0$

$$\pi_+ L_N^{|m|} F_m = -2i(N + |m|)\omega^{\frac{1}{2}} x^{-\frac{1}{2}} L_N^{|m|-1} F_{m+1}, \quad (8)$$

$$\pi_- L_N^{|m|} F_m = 2i\omega^{\frac{1}{2}} x^{\frac{1}{2}} L_N^{|m|+1} F_{m-1}, \quad (9)$$

where m is quantum number. Here [7];

$$N = \frac{1}{4\omega} (E^2 - m^2 - P_z^2) + \frac{(m - |m|)}{2}, \quad \text{for } m > 0, \quad (10)$$

$$N = \frac{1}{4\omega} (E^2 - m^2 - P_z^2) + \frac{(m - |m| - 2)}{2}, \quad \text{for } m < 0 \quad (11)$$

and

$$F_m = \left(\sqrt{\omega\rho^2} \right)^{|m|} e^{im\theta + iP_z z - \frac{x}{2}}, \quad (12)$$

$$\omega = \frac{eB}{2}, \quad x = \omega\rho^2. \quad (13)$$

There are four situations due to spin orientations in magnetic field and positive and negative values of m .

1.Situation: Spin is parallel to the magnetic field and $m > 0$

If we choose the wave function in the form

$$\Psi = \begin{bmatrix} L_N^{|m|} F_m \\ 0 \\ c_1 L_N^{|m|} F_m \\ d_1 L_{N-1}^{|m|+1} F_{m+1} \end{bmatrix} \quad (14)$$

and use it in Eq.(3) we obtain energy spectrum and c_1, d_1 coefficients as follows:

$$E_{\uparrow, m>0}^2 = P_z^2 + m^2 + 4\omega(N-1), \quad (15)$$

$$c_1 = \frac{P_z}{(E+m)}, \quad (16)$$

$$d_1 = \frac{2i\omega^{\frac{1}{2}}x^{\frac{1}{2}}}{(E+m)}. \quad (17)$$

2.Situation: Spin is parallel to the magnetic field and $m<0$

If we choose the wave function in the form

$$\Psi = \begin{bmatrix} L_N^{|m|} F_m \\ 0 \\ c_2 L_N^{|m|} F_m \\ d_2 L_N^{|m|-1} F_{m+1} \end{bmatrix} \quad (18)$$

and use it in Eq.(3) we obtain energy spectrum c_2, d_2 coefficients as follows:

$$E_{\uparrow, m<0}^2 = P_z^2 + m^2 + 4\omega(N+|m|), \quad (19)$$

$$c_2 = \frac{P_z}{(E+m)}, \quad (20)$$

$$d_2 = \frac{-2i(N+|m|)\omega^{\frac{1}{2}}x^{-\frac{1}{2}}}{(E+m)}. \quad (21)$$

3.Situation: Spin is anti-parallel to the magnetic field and $m>0$

If we choose the wave function in the form

$$\Psi = \begin{bmatrix} 0 \\ L_N^{|m|} F_m \\ c_3 L_{N+1}^{|m|-1} F_{m-1} \\ d_3 L_N^{|m|} F_m \end{bmatrix} \quad (22)$$

and use it in Eq.(3) we obtain energy spectrum and c_3, d_3 coefficients as follows:

$$E_{\downarrow, m>0}^2 = P_z^2 + m^2 + 4\omega N, \quad (23)$$

$$c_3 = \frac{-2iN\omega^{\frac{1}{2}}x^{-\frac{1}{2}}}{(E+m)}, \quad (24)$$

$$d_3 = \frac{-P_z}{(E+m)}. \quad (25)$$

4.Situation: Spin is anti-parallel to the magnetic field and $m < 0$

If we choose the wave function in the form

$$\Psi = \begin{bmatrix} 0 \\ L_N^{|m|} F_m \\ c_4 L_N^{m+1} F_{m-1} \\ d_4 L_N^{|m|} F_m \end{bmatrix} \quad (26)$$

and use it in Eq.(3) we obtain energy spectrum and c_4, d_4 coefficients as follows:

$$E_{\downarrow, m < 0}^2 = P_z^2 + m^2 + 4\omega(N + |m| + 1), \quad (27)$$

$$c_4 = \frac{2i\omega^{\frac{1}{2}} x^{\frac{1}{2}}}{(E + m)}, \quad (28)$$

$$d_4 = \frac{-P_z}{(E + m)}. \quad (29)$$

III. EIGENVALUE EQUATION AND ENERGY SPECTRUM OF SPIN-1 PARTICLE

Massive spin-1 particle will be considered as a two-particle system of spin- $\frac{1}{2}$ instead of a single spin-1 particle. This is the second quantization approach of Schrödinger to the problem. For a massive spin-1 particle the Kemmer equation is given in the form

$$(\beta^\mu \pi_\mu - m) \Psi(x) = 0 \quad (30)$$

where 16×16 Kemmer matrices β^μ are given as

$$\beta^\mu = \gamma^\mu \otimes \mathbf{I} + \mathbf{I} \otimes \gamma^\mu$$

with usual γ^μ Dirac matrices. π_μ is electrodynamical four momentum tensor and m is mass of two spin- $\frac{1}{2}$ particles. The geometry of the problem is cylindrically since the particle is rotating around z-axis with a θ angle in xy-plane and going forward to z-axis direction.

The sixteen-component wave function of the equation can be written in the form

$$\Psi = \begin{pmatrix} A \\ B \\ C \\ D \end{pmatrix}$$

where A, B, C, D are 4-component spinors and given as follows:

$$A = \begin{pmatrix} A_+ \\ A_0 \\ A_{\tilde{o}} \\ A_- \end{pmatrix} = \begin{pmatrix} \varphi_+ \varphi_+ \\ \varphi_+ \varphi_- \\ \varphi_- \varphi_+ \\ \varphi_- \varphi_- \end{pmatrix}, B = \begin{pmatrix} B_+ \\ B_0 \\ B_{\tilde{o}} \\ B_- \end{pmatrix} = \begin{pmatrix} \varphi_+ \chi_+ \\ \varphi_+ \chi_- \\ \varphi_- \chi_+ \\ \varphi_- \chi_- \end{pmatrix}$$

$$C = \begin{pmatrix} C_+ \\ C_0 \\ C_{\tilde{o}} \\ C_- \end{pmatrix} = \begin{pmatrix} \chi_+ \varphi_+ \\ \chi_+ \varphi_- \\ \chi_- \varphi_+ \\ \chi_- \varphi_- \end{pmatrix}, D = \begin{pmatrix} D_+ \\ D_0 \\ D_{\tilde{o}} \\ D_- \end{pmatrix} = \begin{pmatrix} \chi_+ \chi_+ \\ \chi_+ \chi_- \\ \chi_- \chi_+ \\ \chi_- \chi_- \end{pmatrix}.$$

In these matrices the left and right elements of the right matrix indicate first and second particles respectively and “+” and “-” indices indicate the spin orientations in magnetic field.

After some algebra Kemmer equation takes the following form

$$\left[\left(\gamma_1^0 \otimes I_2 + I_1 \otimes \gamma_2^0 \right) E + \left(\gamma_1^0 \otimes \vec{\alpha}_2 + \vec{\alpha}_1 \otimes \gamma_2^0 \right) \cdot \vec{\pi} - m \gamma_1^0 \otimes \gamma_2^0 \right] \begin{pmatrix} A \\ B \\ C \\ D \end{pmatrix} = 0 \quad (31)$$

where

$$\vec{\alpha} = \gamma^0 \vec{\gamma} = \begin{pmatrix} 0 & \vec{\sigma} \\ \vec{\sigma} & 0 \end{pmatrix},$$

$$\gamma^0 = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}.$$

We obtain four linear algebraic equations, ultimately from Eq.(31):

$$(2E - m) A - \vec{\sigma}_{(1)} \cdot \vec{\pi} C - \vec{\sigma}_{(2)} \cdot \vec{\pi} B = 0, \quad (32)$$

$$\vec{\sigma}_{(2)} \cdot \vec{\pi} A - \vec{\sigma}_{(1)} \cdot \vec{\pi} D - mB = 0, \quad (33)$$

$$\vec{\sigma}_{(1)} \cdot \vec{\pi} A - \vec{\sigma}_{(2)} \cdot \vec{\pi} D - mC = 0, \quad (34)$$

$$(2E + m) D - \vec{\sigma}_{(1)} \cdot \vec{\pi} B - \vec{\sigma}_{(2)} \cdot \vec{\pi} C = 0, \quad (35)$$

where

$$\vec{\sigma}_{(1)} \cdot \vec{\pi} = (\vec{\sigma} \otimes \mathbf{I}) \cdot \vec{\pi} = \begin{pmatrix} P_z & 0 & \pi_- & 0 \\ 0 & P_z & 0 & \pi_- \\ \pi_+ & 0 & -P_z & 0 \\ 0 & \pi_+ & 0 & -P_z \end{pmatrix}, \quad (36)$$

$$\vec{\sigma}_{(2)} \cdot \vec{\pi} = (\mathbf{I} \otimes \vec{\sigma}) \cdot \vec{\pi} = \begin{pmatrix} P_z & \pi_- & 0 & 0 \\ \pi_+ & -P_z & 0 & 0 \\ 0 & 0 & P_z & \pi_- \\ 0 & \pi_+ & \pi_+ & -P_z \end{pmatrix}, \quad (37)$$

and

$$\pi_+ = \pi_1 + i\pi_2 = e^{i\theta} \left[-i\partial_r + \frac{1}{r}\partial_\theta - \frac{iBre}{2} \right], \quad (38)$$

$$\pi_- = \pi_1 - i\pi_2 = e^{-i\theta} \left[-i\partial_r - \frac{1}{r}\partial_\theta + \frac{iBre}{2} \right]. \quad (39)$$

By using the explicit forms of matrices seen in Eqs.(36), (37) and substituting the following equalities

$$B_0(\varphi_+\chi_-) = C_{\bar{o}}(\chi_-\varphi_+) \quad , B_{\bar{o}}(\varphi_-\chi_+) = C_0(\chi_+\varphi_-),$$

$$A_0(\varphi_+\varphi_-) = A_{\bar{o}}(\varphi_-\varphi_+) \quad , D_0(\chi_+\chi_-) = D_{\bar{o}}(\chi_-\chi_+),$$

into four equations given by Eqs.(32)-(35) we obtain ten linear algebraic equations in the form

$$(2E - m) A_+ - 2P_z B_+ - 2\pi_- B_0 = 0, \quad (40)$$

$$(2E - m) A_0 - \pi_+ B_+ - \pi_- B_- + P_z (B_0 - B_{\bar{0}}) = 0, \quad (41)$$

$$(2E - m) A_- + 2P_z B_- - 2\pi_+ B_{\bar{0}} = 0, \quad (42)$$

$$(2E - m) D_+ - 2P_z B_+ - 2\pi_- B_{\bar{0}} = 0, \quad (43)$$

$$(2E - m) D_0 - \pi_+ B_+ - \pi_- B_- + P_z (B_{\bar{0}} - B_0) = 0, \quad (44)$$

$$(2E - m) D_- + 2P_z B_- - 2\pi_+ B_0 = 0, \quad (45)$$

$$P_z (A_+ - D_+) + \pi_- (A_0 - D_0) - m B_+ = 0, \quad (46)$$

$$P_z (D_- - A_-) + \pi_+ (A_0 - D_0) - m B_- = 0, \quad (47)$$

$$P_z (A_0 + D_0) + \pi_- A_- - \pi_+ D_+ - m B_{\bar{0}} = 0, \quad (48)$$

$$P_z (A_0 + D_0) - \pi_+ A_+ + \pi_- D_- + m B_0 = 0. \quad (49)$$

In order to find the energy eigenvalues we will choose B_0 and $B_{\bar{0}}$ in terms of Laguerre polynomials:

$$B_0 = L_N^{[m]} F_m,$$

$$B_{\bar{0}} = -L_N^{[m]} F_m.$$

(i) for $m > 0$

If we use $B_0 = L_N^{[m]} F_m$ and $B_{\bar{0}} = -L_N^{[m]} F_m$ in Eqs.(40)-(49) we find A, B, C, D spinors in the form

$$B = \begin{pmatrix} b_+ \omega^{\frac{1}{2}} x^{-\frac{1}{2}} L_{N+1}^{[m]-1} F_{m-1} \\ L_N^{[m]} F_m \\ -L_N^{[m]} F_m \\ b_- \omega^{\frac{1}{2}} x^{\frac{1}{2}} L_{N-1}^{[m]+1} F_{m+1} \end{pmatrix}, \quad C = \begin{pmatrix} b_+ \omega^{\frac{1}{2}} x^{-\frac{1}{2}} L_{N+1}^{[m]-1} F_{m-1} \\ -L_N^{[m]} F_m \\ L_N^{[m]} F_m \\ b_- \omega^{\frac{1}{2}} x^{\frac{1}{2}} L_{N-1}^{[m]+1} F_{m+1} \end{pmatrix},$$

$$A = \begin{pmatrix} \frac{2}{2E-m} (P_z b_+ - 2iN) \omega^{\frac{1}{2}} x^{-\frac{1}{2}} L_{N+1}^{[m]-1} F_{m-1} \\ \frac{1}{2E-m} [2i\omega (b_+ - (N-1)b_-) - 2P_z] L_N^{[m]} F_m \\ \frac{1}{2E-m} [2i\omega (b_+ - (N-1)b_-) - 2P_z] L_N^{[m]} F_m \\ -\frac{2}{2E-m} (P_z b_- + 2i) \omega^{\frac{1}{2}} x^{\frac{1}{2}} L_{N-1}^{[m]+1} F_{m+1} \end{pmatrix},$$

$$D = \begin{pmatrix} \frac{2}{2E+m} (P_z b_+ + 2iN) \omega^{\frac{1}{2}} x^{-\frac{1}{2}} L_{N+1}^{|m|-1} F_{m-1} \\ \frac{1}{2E+m} [2i\omega (b_+ - (N-1)b_-) + 2P_z] L_N^{|m|} F_m \\ \frac{1}{2E+m} [2i\omega (b_+ - (N-1)b_-) + 2P_z] L_N^{|m|} F_m \\ -\frac{2}{2E+m} (P_z b_- - 2i) \omega^{\frac{1}{2}} x^{\frac{1}{2}} L_{N-1}^{|m|+1} F_{m+1} \end{pmatrix}.$$

We obtain four algebraic equations by using these spinors in Eqs.(46)-(49) in the form

$$\left[E^2 - P_z^2 - \frac{m^2}{4} - 2\omega N \right] b_+ + 2N(N-1)\omega b_- = 0, \quad (50)$$

$$\left[E^2 - P_z^2 - \frac{m^2}{4} - 2\omega(N-1) \right] b_- + 2\omega b_+ = 0, \quad (51)$$

$$\left[E^2 - P_z^2 - \frac{m^2}{4} + 4\omega \left(\frac{E}{m} - \left(N - \frac{1}{2} \right) \right) \right] + i\omega P_z (N-1) b_- + i\omega P_z b_+ = 0, \quad (52)$$

$$\left[E^2 - P_z^2 - \frac{m^2}{4} - 4\omega \left(\frac{E}{m} + \left(N - \frac{1}{2} \right) \right) \right] - i\omega P_z (N-1) b_- - i\omega P_z b_+ = 0. \quad (53)$$

From Eqs.(50)-(52) we find

$$b_+ = \frac{2N\omega \left[E^2 - P_z^2 - \frac{m^2}{4} + 4\omega \left(\frac{E}{m} - \left(N - \frac{1}{2} \right) \right) \right]}{i\omega P_z \left[E^2 - P_z^2 - \frac{m^2}{4} - 4\omega N \right]}, \quad (54)$$

$$b_- = \frac{- \left[E^2 - P_z^2 - \frac{m^2}{4} + 4\omega \left(\frac{E}{m} - \left(N - \frac{1}{2} \right) \right) \right] \left[E^2 - P_z^2 - \frac{m^2}{4} - 2\omega N \right]}{i\omega P_z (N-1) \left[E^2 - P_z^2 - \frac{m^2}{4} - 4\omega N \right]}. \quad (55)$$

If we substitute these into Eqs.(51)-(53) we obtain energy levels as follows

$$E^2 = P_z^2 + \frac{m^2}{4} + 4\omega \left(N - \frac{1}{2} \right), \quad (56)$$

$$E = \frac{2\omega}{m} \pm \frac{1}{2} \sqrt{\frac{16\omega^2}{m^2} + m^2 + 4P_z^2 + 16\omega \left(N - \frac{1}{2} \right)}, \quad (57)$$

$$E = -\frac{2\omega}{m} \pm \frac{1}{2} \sqrt{\frac{16\omega^2}{m^2} + m^2 + 4P_z^2 + 16\omega \left(N - \frac{1}{2} \right)}. \quad (58)$$

(ii) for $m < 0$

If we use $B_0 = L_N^{|m|} F_m$ and $B_{\bar{0}} = -L_N^{|m|} F_m$ in Eqs.(40)-(49) we find A, B, C, D spinors in the form

$$B = \begin{pmatrix} b_+ \omega^{\frac{1}{2}} x^{\frac{1}{2}} L_N^{|m|+1} F_{m-1} \\ L_N^{|m|} F_m \\ -L_N^{|m|} F_m \\ b_- \omega^{\frac{1}{2}} x^{-\frac{1}{2}} L_N^{|m|-1} F_{m+1} \end{pmatrix}, \quad C = \begin{pmatrix} b_+ \omega^{\frac{1}{2}} x^{\frac{1}{2}} L_N^{|m|+1} F_{m-1} \\ -L_N^{|m|} F_m \\ L_N^{|m|} F_m \\ b_- \omega^{\frac{1}{2}} x^{-\frac{1}{2}} L_N^{|m|-1} F_{m+1} \end{pmatrix},$$

$$A = \begin{pmatrix} \frac{2}{2E-m} (P_z b_+ + 2i) \omega^{\frac{1}{2}} x^{\frac{1}{2}} L_N^{|m|+1} F_{m-1} \\ \frac{1}{2E-m} [-2i\omega (N + |m| + 1) b_+ + 2i\omega b_- - 2P_z] L_N^{|m|} F_m \\ \frac{1}{2E-m} [-2i\omega (N + |m| + 1) b_+ + 2i\omega b_- - 2P_z] L_N^{|m|} F_m \\ \frac{2}{2E-m} [-P_z b_- + 2i (N + |m|)] \omega^{\frac{1}{2}} x^{-\frac{1}{2}} L_N^{|m|-1} F_{m+1} \end{pmatrix},$$

$$D = \begin{pmatrix} \frac{2}{2E-m} (P_z b_+ - 2i) \omega^{\frac{1}{2}} x^{\frac{1}{2}} L_N^{|m|+1} F_{m-1} \\ \frac{1}{2E-m} [-2i\omega (N + |m| + 1) b_+ + 2i\omega b_- + 2P_z] L_N^{|m|} F_m \\ \frac{1}{2E-m} [-2i\omega (N + |m| + 1) b_+ + 2i\omega b_- + 2P_z] L_N^{|m|} F_m \\ -\frac{2}{2E-m} [P_z b_- + 2i (N + |m|)] \omega^{\frac{1}{2}} x^{-\frac{1}{2}} L_N^{|m|-1} F_{m+1} \end{pmatrix}.$$

We obtain four algebraic equations by using these spinors in Eqs.(50)-(??) in the form

$$\left[E^2 - P_z^2 - \frac{m^2}{4} - 2\omega (N + |m| + 1) \right] b_+ + 2\omega b_- = 0 \quad (59)$$

$$\left[E^2 - P_z^2 - \frac{m^2}{4} + 4\omega \left(\frac{E}{m} - (N + |m| + \frac{1}{2}) \right) \right] - i\omega P_z b_- - i\omega \left(N + |m| + \frac{1}{2} \right) P_z b_+ = 0 \quad (60)$$

$$\left[E^2 - P_z^2 - \frac{m^2}{4} - 4\omega \left(\frac{E}{m} + (N + |m| + \frac{1}{2}) \right) \right] + i\omega P_z b_- + i\omega \left(N + |m| + \frac{1}{2} \right) P_z b_+ = 0 \quad (61)$$

$$\left[E^2 - P_z^2 - \frac{m^2}{4} - 2\omega (N + |m|) \right] b_- + 2\omega (N + |m|) (N + |m| + 1) b_+ = 0 \quad (62)$$

From Eqs.(50)-(52) we find

$$b_+ = \frac{2\omega \left[E^2 - P_z^2 - \frac{m^2}{4} - 4\omega \left(\frac{E}{m} + (N + |m| + \frac{1}{2}) \right) \right]}{i\omega P_z \left[E^2 - P_z^2 - \frac{m^2}{4} - 4\omega (N + |m| + 1) \right]}, \quad (63)$$

$$b_- = \frac{- \left[E^2 - P_z^2 - \frac{m^2}{4} - 4\omega \left(\frac{E}{m} + (N + |m| + \frac{1}{2}) \right) \right] \left[E^2 - P_z^2 - \frac{m^2}{4} - 2\omega (N + |m| + 1) \right]}{i\omega P_z \left[E^2 - P_z^2 - \frac{m^2}{4} - 4\omega (N + |m| + 1) \right]}, \quad (64)$$

If we substitute these into Eqs.(51)-(53) we obtain energy levels as follows

$$E^2 = P_z^2 + \frac{m^2}{4} + 4\omega \left(N + |m| + \frac{1}{2} \right), \quad (65)$$

$$E = \frac{2\omega}{m} \pm \frac{1}{2} \sqrt{\frac{16\omega^2}{m^2} + m^2 + 4P_z^2 + 16\omega \left(N + |m| + \frac{1}{2} \right)}, \quad (66)$$

$$E = -\frac{2\omega}{m} \pm \frac{1}{2} \sqrt{\frac{16\omega^2}{m^2} + m^2 + 4P_z^2 + 16\omega \left(N + |m| + \frac{1}{2} \right)}. \quad (67)$$

IV. CONCLUSIONS

There are a few techniques developed for understanding the behavior of spin-1 particles moving in a constant magnetic field. These techniques are quite complicated. Because of no available experimental tests of these techniques we do not know which is the best.

We used here a simple algebraic method for obtaining the energy levels of spin-1 particle moving in a constant magnetic field. Spin-1 particle was considered as a two-fermion system and in the process to obtain the energy levels the wave functions of spin- $\frac{1}{2}$ particle which are written in terms of Laguerre polynomials were used.

If the magnetic field is quite weak, we see that the difference of the second and third from the first energy level is $\pm\frac{2\omega}{m}$. This comes from the spin orientations in magnetic field. The second and third energy spectrums can be interpreted as the energy spectrums of situations in which the spins of particle and anti-particle are parallel and anti-parallel to the magnetic field, respectively. From these spectrums it is seen for all values of magnetic field the transition energy of spin from $s=1$ to $s=-1$ is equal to

$$\Delta E_{\pm} = \frac{4\omega}{m} \quad .$$

REFERENCES

- [1] N. Kemmer, Proc. Soc. (London) **A 173**, 91 (1939).
- [2] D. Shay, R. H. Good, Phys. Rev. **179**, 1410 (1969).
- [3] W.Tsai, Phys. Rev. **D 4(12)**, 3652 (1971).
- [4] L. D. Krase, P. Lu, R. H. Good, Phys. Rev. **D 3(6)**, 1275 (1971).
- [5] W. Tsai and A.Yildiz, Phys. Rev. **D 4**, 3643 (1971).
- [6] F. Halzen and A. D. Martin, *Quarks and Leptons:An Introductory Course in Modern Particle Physics*, J. Wiley and Sons, Inc. New York, (1984).
- [7] A. Havare and F. Binbay, Int. J. Diff. Eq. and App., **1 (3)**, (2000).